



Reference values for serum leptin in healthy non-obese children and adolescents

Ulrik Lausten-Thomsen, Michael Christiansen, Paula Louise Hedley, Cilius Esmann Fonvig, Theresa Stjernholm, Oluf Pedersen, Torben Hansen & Jens-Christian Holm

To cite this article: Ulrik Lausten-Thomsen, Michael Christiansen, Paula Louise Hedley, Cilius Esmann Fonvig, Theresa Stjernholm, Oluf Pedersen, Torben Hansen & Jens-Christian Holm (2016) Reference values for serum leptin in healthy non-obese children and adolescents, *Scandinavian Journal of Clinical and Laboratory Investigation*, 76:7, 561-567, DOI: [10.1080/00365513.2016.1210226](https://doi.org/10.1080/00365513.2016.1210226)

To link to this article: <https://doi.org/10.1080/00365513.2016.1210226>



Published online: 09 Aug 2016.



Submit your article to this journal [↗](#)



Article views: 241



View Crossmark data [↗](#)



Citing articles: 3 View citing articles [↗](#)

ORIGINAL ARTICLE

Reference values for serum leptin in healthy non-obese children and adolescents

Ulrik Lausten-Thomsen^{a,b}, Michael Christiansen^{c,d}, Paula Louise Hedley^c, Cilius Esmann Fonvig^{a,e},
Theresa Stjernholm^a, Oluf Pedersen^e, Torben Hansen^{e,f} and Jens-Christian Holm^{a,b,e}

^aDepartment of Pediatrics, The Children's Obesity Clinic, Copenhagen University Hospital Holbæk, Holbæk, Denmark; ^bDepartment of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark; ^cDepartment for Congenital Disorders, Statens Serum Institut, Copenhagen, Denmark; ^dDepartment of Biomedical Sciences, University of Copenhagen, Copenhagen, Denmark; ^eThe Faculty of Health and Medical Sciences, Section of Metabolic Genetics, The Novo Nordisk Foundation Center for Basic Metabolic Research, University of Copenhagen, Copenhagen, Denmark; ^fFaculty of Health Sciences, University of Southern Denmark, Odense, Denmark

ABSTRACT

Background: Adipokines are biologically active, low-molecular weight peptides, which play a major role in metabolic homeostasis in humans. Leptin has gained increasing attention in pediatrics as a biomarker for various metabolic pathologies. Yet, its usefulness is hampered by the relative lack of reference values from pediatric settings. Accordingly, this study aims to evaluate serum concentrations of leptin, soluble leptin receptor (sOB-R), and free leptin index (FLI) in healthy Danish schoolchildren aged 6–18 years and subsequently to establish reference intervals across sex and age groups.

Methods: A total of 1193 healthy, non-obese Danish schoolchildren (730 girls, 463 boys) aged 6–18 years (median 11.9) were examined by trained medical staff. Serum leptin and sOB-R concentrations in venous fasting blood samples were quantitated by immunoassay. Percentile curves of leptin, sOB-R, and free leptin index were calculated using the General Additive Model for Location Scale and Shape (GAMLSS).

Results: Significant age and sex-dependent differences in circulating leptin levels were found. In boys, the median leptin concentration for all ages combined was 3.35 µg/L (95%-interval: 0.71–22.47) and in girls, it was 9.89 ng/L (95%-interval: 2.06–41.49). For sOB-R, no sex-specific difference was found, and the median sOB-R concentration was 8.24 µg/L (IQR: 3.58–23.74; range: < 1.56–744.15).

Conclusion: We demonstrated an age-dependent correlation with both serum leptin concentration and free leptin index with a gradual and significant increase in girls throughout childhood and adolescence and a significantly higher leptin concentration and free leptin index bell-shaped peak in early adolescence in boys.

ARTICLE HISTORY

Received 26 May 2016
Revised 23 June 2016
Accepted 4 July 2016
Published online 2 August 2016

KEYWORDS

Adipokines; adolescents; child; humans; leptin receptors; leptin; reference values

Introduction

Adipose tissue is an important endocrine organ which secretes several biologically active, low-molecular weight peptides, or adipokines, which play a major role in human metabolic homeostasis [1]. Leptin is an important adipokine that is synthesized mainly, but not exclusively, by white adipose tissue and has vast endocrine and metabolic functions [1,2]. Leptin is involved in the regulation of feeding and energy balance [3,4] and serves as a mediator of the neuroendocrinological adaptation to fasting by influencing the expression of different neuropeptides that regulates energy expenditure [2,5]. Furthermore, leptin is implicated in fatty acid metabolism [6], in the reproductive system [7,8], in the immune system [9–11], and the serum leptin concentration has been proposed as a predictor of metabolic risks such as insulin resistance [12]. Accordingly, leptin is increased in obese males and females compared to non-obese individuals [13,14].

Importantly, the bioavailability of leptin is modulated by the soluble leptin receptor (sOB-R) [15]. Leptin achieves signal transduction via four different receptor (OB-R) isoforms,

but the amount of functionally active OB-R is affected by constitutive shedding of the extra-cellular domain [16]. The product of the cleavage process, i.e. the sOB-R, is the main binding protein for leptin [17] and thus modulates its bioavailability [18]. It has been proposed that the ratio between leptin and sOB-R, or the free leptin index (FLI), may be a useful tool to assess leptin activity [19].

Leptin has gained increasing attention in pediatrics as a serum biomarker due to its correlation to various metabolic risk factors such as insulin resistance, metabolic syndrome, and cardiovascular disease [20–22]. Childhood obesity has increased dramatically across ethnic groups in recent decades, and with that the obesity-associated comorbidities have risen in importance [23]. Consequently, the need for cardio-metabolic risk stratification has increased and created a demand for pediatric reference values for relevant biomarkers, including leptin, the sOB-R, and their mutual relationship.

However, the interpretation of serum values in children, and thus an assessment of the clinical utility of the analyte, is hampered by the relative lack of reference values from a pediatric setting. Contrary to adults, only a few studies on

children and adolescents have been published [24]. Furthermore, the available data is largely obtained in small and heterogeneous study materials containing a limited number of healthy children [25].

The objective of the present study is to elucidate the naturally occurring serum concentrations of leptin and sOB-R and calculate the FLI in a large population-based sampled cohort of healthy, non-obese Danish schoolchildren and furthermore, to establish reference intervals for children and adolescents across sex and age groups.

Material

Subjects

Danish schoolchildren from several municipalities in the region of Zealand, Denmark, were included between October 2010 and November 2013. Exclusion criteria were obesity defined as a body mass index (BMI) above the 95th percentile for sex and age [26] at the time of inclusion, and known diseases requiring regular medication. All participants and/or parents signed informed consent. The study was approved by the Danish Data Protection Agency (REG-06-2014) and the Regional Scientific Ethics Committee (protocol no. SJ-104).

Anthropometric measurements

The participants were examined by skilled research assistants immediately prior to blood sampling. Children's heights were measured using a stadiometer to the nearest millimeter with the subject standing without shoes. The weight was measured using a BC-418 Segmental Body Composition Analyzer (Tanita, Tokyo, Japan) to the nearest 100 grams; with subjects wearing light indoor clothes and no shoes. The degree of baseline BMI is expressed in standard deviation score (SDS) in order to adjust for age- and sex-related variation in reference intervals. BMI SDS was calculated by the LMS method by converting BMI into a normal distribution by sex and age using the median coefficient of variation and a measure of the skewness [27] based on the Box-Cox power plot from Danish BMI charts [28].

Additional data collection

Additional data on health and socioeconomic status were obtained through a structured family-based questionnaire. Similarly, the pubertal developmental stage was assessed by a questionnaire where standardized pictures of the five pubertal developmental stages suggested by Marshall and Tanner [29,30] are depicted accompanied by a text describing each category and thereby allowing the participants and their parents in estimating the appropriate pubertal stage *ad modum* Tanner

Biochemical analyses

After an overnight fast, venous blood samples were collected from each child between 07:00 and 09:00 h. The samples were processed immediately and the serum was stored at

−80 °C until further analysis. Serum leptin concentration was quantitated *in singlo* using an optimized version of the Human Leptin ELISA Development Kit, DuoSet (DY398, R&D Systems, Minneapolis, MN, USA) following appropriate sample dilution. The detection limit of the assay was 0.0312 µg/L. Assay calibrator was highly purified *Escherichia coli* expressed recombinant human leptin produced by the manufacturer (AF398, R&D Systems, Minneapolis, MN, USA).

The sOB-R was quantitated *in singlo* using the Quantikine Human Leptin sR Immunoassay (DOBR00, R&D Systems, Minneapolis, MN, USA) as described by the manufacturer following appropriate sample dilution. The smallest sOB-R concentration detectable within a linear standard curve was 1.56 µg/L when the serum was diluted 1:4. When standards were diluted in serum to extend the standard curve, the dilution curve was not linear, suggesting a matrix-effect on the measurements. Assay calibrator was highly purified NS0-expressed recombinant leptin R/Fc chimera produced by the manufacturer.

For both leptin and sOB-R, the intra- and inter-assay coefficients of variations were <5% and <10%, respectively. At room temperature, leptin concentrations in serum were stable for 48 h, and sOB-R was stable for 56 h. Both analytes were stable for three months at −20 °C, and for 10 freeze-thaw cycles.

Statistical analyses


Statistical modelling was performed using the statistical software R, version 3.2.4 (R Foundation for Statistical Computing, Vienna, Austria). Skewness was estimated by D'Agostino test and log transformation of data was performed when appropriate. Differences between sexes were evaluated non-parametrically for the entire group and for each age interval using the Wilcoxon-Mann-Whitney test. Similarly, Wilcoxon-Mann-Whitney tests were used to compare specific age groups to the defined reference group of 16–18-year-old adolescents. As studies have demonstrated that healthy girls experience a graduate increase in body fat percentage throughout childhood and adolescence whereas healthy boys have a more stable body composition and often even decreasing fat percentage [31], the data was analyzed separately for each sex and subsequently adjusted for age and BMI SDS. As concentrations of sOB-R can provide an indication of free leptin, the free leptin index (FLI) being defined as the ratio of leptin to sOB-R [32] was calculated as $[\text{leptin}]/[\text{sOB-R}] \times 100$.

Percentile curves of leptin, sOB-R, and FLI were calculated as a function of age as a continuous variable using the General Additive Model for Location Scale and Shape (GAMLSS) [33]. Percentiles for age groups applied exactly for the midpoint of each age group were equally calculated with GAMLSS. Furthermore, the correlation between leptin, sOB-R, and FLI and age was evaluated using a generalized linear model (GLM), adjusted for BMI SDS.

To avoid left censoring of data in cases of values below the limit of detection (LOD), we used a tobit regression (TOBIT) model [34] instead of the more traditional approach of substitution method of replacing the values below LOD by

Table 1. Percentiles of leptin ($\mu\text{g/L}$) calculated with the General Additive Model for Location Scale and Shape (GAMLSS).

Age (years)	Percentiles for girls							Percentiles for boys						
	2.5	5	25	50	75	95	97.5	2.5	5	25	50	75	95	97.5
6	0.92	1.19	2.53	4.18	6.88	14.45	18.61	0.44	0.55	1.15	1.97	3.47	8.22	11.03
7	1.09	1.40	2.93	4.80	7.83	16.09	20.52	0.54	0.68	1.37	2.29	3.94	9.03	11.99
8	1.29	1.65	3.40	5.52	8.93	17.99	22.75	0.68	0.84	1.65	2.72	4.59	10.25	13.53
9	1.54	1.95	3.96	6.37	10.21	20.23	25.37	0.83	1.02	1.99	3.26	5.49	12.30	16.23
10	1.82	2.30	4.60	7.34	11.65	22.71	28.29	0.99	1.23	2.41	3.99	6.82	15.70	20.96
11	2.14	2.69	5.32	8.42	13.26	25.45	31.51	1.14	1.41	2.84	4.81	8.45	20.54	28.03
12	2.54	3.18	6.21	9.76	15.24	28.85	35.51	1.10	1.38	2.88	5.04	9.21	24.11	33.91
13	3.04	3.79	7.32	11.43	17.71	33.06	40.48	0.88	1.11	2.40	4.35	8.32	23.80	34.73
14	3.58	4.44	8.49	13.16	20.24	37.30	45.45	0.66	0.84	1.86	3.44	6.75	20.53	30.78
15	4.07	5.04	9.53	14.66	22.38	40.74	49.42	0.55	0.69	1.51	2.77	5.41	16.45	24.73
16	4.46	5.50	10.30	15.73	23.84	42.90	51.81	0.51	0.64	1.34	2.37	4.45	12.65	18.57
17	4.71	5.79	10.75	16.31	24.53	43.65	52.50	0.52	0.64	1.24	2.07	3.63	9.16	12.81
18	4.84	5.94	10.92	16.46	24.59	43.27	51.84	0.53	0.63	1.13	1.77	2.88	6.33	8.41

**Figure 1.** (a) Leptin (y-axis) in healthy girls as function of age (x-axis). The 97.5th, 95th, 50th, and 5th percentiles calculated with GAMLSS are presented. (b) Leptin (y-axis) in healthy boys as function of age (x-axis). The 97.5th, 95th, 50th, and 5th percentiles calculated with GAMLSS are presented.

a fixed value [35] as the TOBIT approach has been reported to perform better when the proportion of non-detects, i.e. values below LOD, is below 30% [36].

Results

Included patients

A total of 1196 healthy children (730 girls and 463 boys) aged 6–18 (median 11.9) years old were included in the study. The children had normal weight (median 42.5 kg, range 18.4–85.7) for height (median 154.0 cm, range: 111.6–195.0) conveying a median BMI SDS of 0.34 (range: –1.65 to +1.65). There were no statistically significant associations with social class or BMI SDS between sex and age groups (data not shown).

Leptin

The combined leptin concentration for all included children was non-normally distributed (Shapiro-Wilk test, $W = 0.7729$, $p < 2.2e-16$) with a combined median for both genders and all ages of 7.06 $\mu\text{g/L}$ (IQR: 3.12–13.49; 95%-interval: 0.87–29.56; range: 0.45–89.01). The leptin concentration was found to be

significantly elevated in girls when compared to boys (Mann-Witney Wilcoxon U test, $p < 0.0001$) throughout the age groups, and data was subsequently analyzed separately for each sex.

In girls, the median leptin concentration for all ages combined was 9.89 $\mu\text{g/L}$ (IQR: 5.69–16.92; 95%-interval: 2.06–41.49; range 0.55–89.01) and in boys, it was 3.35 $\mu\text{g/L}$ (IQR: 1.90–6.42; 95%-interval: 0.71–22.47; range 0.45–60.19).

The calculation of reference intervals was performed in a GAMLSS model using age as a continuous variable and is presented in Table 1 and Figure 1(a,b).

In a generalized linear model, the total serum leptin concentration was correlated to age in girls ($p < 0.0001$), as leptin concentrations increased with age. For boys, serum leptin concentration exhibited a bell-shaped correlation to age with a maximum at 12.5 years as demonstrated by a generalized 2nd order polynomial regression model ($p < 0.0001$). When adjusted for BMI SDS, i.e. the degree of obesity, this age-dependent association persisted in both girls ($p < 0.0001$) and boys ($p < 0.0001$).

sOB-R

The combined median sOB-R concentration for all included children was 8.24 $\mu\text{g/L}$ (IQR: 3.58–23.74; range: <1.56–744.15).

Table 2. Percentiles of sOB-R ($\mu\text{g/L}$) calculated with General Additive Model for Location Scale and Shape (GAMLSS). Values $<1.56 \mu\text{g/L}$ are derived from TOBIT regression. See text for details.

Age (years)	Percentiles for girls							Percentiles for boys						
	2.5	5	25	50	75	95	97.5	2.5	5	25	50	75	95	97.5
6	1.02	1.62	5.46	12.61	36.22	408.65	1229.99	1.74	2.11	4.46	9.46	27.87	332.50	995.86
7	1.70	2.25	5.56	12.03	34.57	402.03	1217.60	1.65	2.00	4.28	9.14	27.23	335.43	1021.19
8	1.72	2.19	5.08	10.83	31.12	364.33	1105.25	1.24	1.60	3.85	8.49	25.47	317.08	975.52
9	1.52	1.94	4.49	9.55	27.39	319.94	970.56	1.14	1.49	3.67	8.14	24.49	308.26	955.57
10	1.33	1.71	3.97	8.44	24.14	281.16	852.83	1.43	1.75	3.80	8.21	24.81	321.44	1007.02
11	1.31	1.63	3.63	7.64	21.85	255.65	776.46	1.26	1.55	3.49	7.80	24.79	369.31	1242.48
12	1.30	1.59	3.38	7.04	20.13	236.81	720.27	1.26	1.55	3.49	7.80	24.79	369.31	1242.48
13	1.25	1.51	3.15	6.53	18.66	219.77	668.90	1.17	1.45	3.36	7.73	25.78	438.62	1578.37
14	1.18	1.43	2.96	6.11	17.44	205.07	624.30	1.09	1.36	3.22	7.61	26.41	502.14	1916.26
15	1.12	1.35	2.80	5.78	16.44	192.97	587.5	1.05	1.31	3.11	7.33	25.30	479.01	1830.05
16	1.07	1.29	2.67	5.50	15.63	183.04	557.32	1.07	1.32	3.02	6.90	22.73	383.19	1384.66
17	1.03	1.24	2.56	5.28	14.95	174.65	531.79	1.10	1.35	2.94	6.42	19.80	282.02	938.83
18	1.00	1.20	2.47	5.08	14.36	167.28	509.35	1.14	1.38	2.87	5.97	17.20	206.07	631.01

A total of 115 children (9.6%) had values below the LOD of $1.56 \mu\text{g/L}$. An attempt to further expand the lower detection limit though dilutions of the calibrator proved futile due to the matrix effect [37,38]. An estimation of the values was calculated using TOBIT regression. The data was non-normally distributed (Shapiro-Wilk test, $W = 0.34$, $p < 2.2e-16$). No difference in sOB-R concentrations between sexes was demonstrated when using a non-parametric Wilcoxon-Mann-Whitney test ($p = 0.47$), a finding which is in accordance with a previous report by Kratzsch et al. [19].

The calculation of reference intervals was performed in a GAMLSS model using age as a continuous variable and is presented in Table 2.

In a generalized linear model, no statistic indication of age-dependent correlation was found in sOB-R, regardless of adjustment for BMI SDS, in girls ($p = 0.21$) or boys ($p = 0.81$). A non-parametric test (Spearman's rank correlation coefficient) demonstrated no correlation between age and sOB-R in girls ($p = 0.57$) or boys ($p = 0.37$).

The sub-population of children with sOB-R values below LOD did not differ from the rest of the population in regards to sex, social class, or BMI SDS (Wilcoxon-Mann-Whitney test; $p = 0.87$, $p = 0.60$, and $p = 0.98$, respectively), but had a higher median age ($p = 0.0001$) and leptin concentration ($p = 0.0152$).

Free-leptin index, FLI

The empirically observed median values for leptin concentration and FLI were calculated independently in each age-group and their course was found to be correlated for both girls and boys (Spearman's rank correlation coefficient, $\rho = 0.49$, $p < 0.0001$ and $\rho = 0.48$, $p < 0.0001$, respectively) and is presented in Figure 2(a,b). The calculation of smoothed reference intervals was performed in a GAMLSS model using age as a continuous variable and is presented in Table 3.

In girls, both leptin concentration and FLI increased significantly during childhood and adolescence and an adjusted GLM model demonstrated a relationship between age and FLI ($p < 0.0001$). Similarly, comparison of individual age groups to late adolescence, i.e. ages 16–18, demonstrated significantly lower FLI values in the younger age groups (i.e. ages 6–10, $p < 0.0001$; ages 11 and 12, $p < 0.01$; and ages 13, $p < 0.05$; Wilcoxon-Mann-Whitney) when compared to

late adolescence, i.e. ages 16–18. In boys however, a GLM model for all ages, showed no significant correlation to age when observing all age groups ($p = 0.947$), but were present when observing boys < 14 years ($p < 0.01$). Accordingly, both leptin concentrations and FLI in boys were significantly higher during early adolescence (i.e. ages 10, 11, and 12; $p = 0.049$, $p = 0.014$, and $p = 0.005$, respectively) when compared by a Wilcoxon-Mann-Whitney test to late adolescence, i.e. ages 16–18.

Relation with pubertal stage

Of the 1196 included children, the pubertal stages were obtained in a subset of 747 cases (495 girls, 252 boys). When using an adjusted GLM model to analyze the relationship between Tanner stages and leptin, sOB-R, and FLI, there was no continuous relationship with sOB-R in neither girls nor boys ($p = 0.363$ and $p = 0.178$, respectively), whereas both leptin ($p < 0.0001$ and $p = 0.0003$, respectively) and FLI ($p < 0.0001$ and $p = 0.0043$, respectively) was significantly correlated to Tanner stages.

For FLI, we performed a selective analysis of levels of the individual pubertal stages in girls and boys. We found significantly elevated sOB-R concentrations in Tanner stage I in girls ($p < 0.0001$) but no difference in boys ($p > 0.05$), when comparing to Tanner stage V by Wilcoxon-Mann-Whitney tests.

Discussion

Pediatric reference intervals

Many laboratory variables in pediatrics vary with sex, age, growth, and development and consequently the reference values for a given variable should ideally be established in the pediatric age ranges for both sexes. Various methods for establishing reference intervals have been proposed [39]. However, practical and ethical considerations often preclude the blood sampling in healthy children, which reduces the possibility of collecting large samples from a standard population of 'normal' children. Accordingly, the data on normative values for leptin in children are mainly based on smaller studies [24,25]. It is well-known that preanalytical, analytical, and postanalytical sources of variation can significantly affect the measurement of leptin concentration in the blood which makes direct




Figure 2. (a) Relationship between leptin and FFI in girls. Empiric values for 50th percentile for leptin (squares) and FFI (triangles) for each age group. (b) Relationship between leptin and FFI in boys. Empiric values for 50th percentile for leptin (squares) and FFI (triangles) for each age group.

Table 3. Percentiles of FFI calculated with GAMLSS.

Age (years)	Percentiles for girls							Percentiles for boys						
	2.5	5	25	50	75	95	97.5	2.5	5	25	50	75	95	97.5
6	0.71	1.38	8.31	24.23	63.32	216.45	312.36	0.27	0.63	5.10	16.45	45.21	157.36	226.35
7	0.82	1.63	10.23	30.59	81.56	285.40	414.55	0.37	0.86	6.95	22.40	61.58	214.61	308.87
8	0.96	1.93	12.69	38.90	105.93	380.37	556.64	0.51	1.16	9.34	30.10	82.75	288.79	415.86
9	1.11	2.29	15.80	49.67	138.20	509.11	750.49	0.67	1.52	12.22	39.34	108.17	377.97	544.58
10	1.26	2.66	19.50	63.05	179.01	672.91	996.80	0.85	1.94	15.53	49.96	137.41	480.76	693.09
11	1.31	2.91	23.44	78.31	226.36	858.91	1272.64	1.06	2.40	19.14	61.55	169.31	593.17	855.61
12	1.22	2.93	26.90	93.38	274.03	1038.02	1531.28	1.27	2.86	22.72	72.97	200.80	704.39	1016.62
13	0.99	2.68	29.58	107.35	318.93	1192.99	1744.72	1.45	3.26	25.75	82.65	227.50	799.10	1153.96
14	0.65	2.17	31.61	120.98	362.23	1319.13	1902.53	1.59	3.56	27.96	89.69	246.92	868.45	1254.82
15	0.34	1.56	33.38	135.57	405.86	1415.01	2001.55	1.69	3.78	29.56	94.74	260.91	918.87	1328.45
16	0.19	1.16	35.90	152.67	451.08	1484.50	2052.60	1.78	3.96	30.82	98.73	271.95	959.01	1387.25
17	0.17	1.08	40.05	172.14	493.91	1521.73	2055.71	1.85	4.12	31.86	101.98	280.98	992.14	1435.98
18	0.21	1.28	45.85	192.02	528.91	1525.15	2016.67	1.91	4.24	32.77	104.81	288.86	1021.25	1478.93

comparison of various studies difficult [25]; yet there is a need to establish reference values for leptin concentrations in children [24].

Our study

In this study, the serum concentrations of leptin and sOB-R in a large cohort of 1196 healthy, non-obese Danish schoolchildren were investigated. The data fills a gap in the literature, as normative data on serum leptin concentrations and particularly on sOB-R in children are sparse when compared to adults [24,25]. Furthermore, very little of the available data are derived from healthy, non-obese children and accordingly many conveniently available pediatric study materials that have been used as reference groups may occasionally harbor biases, that should be identified and adjusted. In contrast, the group of children that was examined in the present study is less likely to be biased as the children were unselected healthy schoolchildren.

Leptin in relation to previous publications

We demonstrated an age-dependent correlation with both serum leptin concentration and FFI with a gradual increase in girls throughout childhood and adolescence. In contrast, boys had significantly higher leptin concentration and FFI in early adolescence. Both these findings are in accordance with previously published data [19,24,40]; albeit our data did not study pre-school children which hinder direct comparison. Still, we demonstrated relatively high levels of leptin in boys

at the onset of puberty which is in concordance with previous finding [19,41], and it is believed to reflect the influence of rapidly rising serum sex steroids concentration on leptin production [42].

Finally, the observed difference in leptin concentrations between our data and previous studies may reflect genetic and/or metabolic differences in the study populations. However, it may also reflect technical differences such as the known difficulties in leptin measurements [25] and it has long been stated that caution should be taken when comparing absolute leptin concentrations across studies [43]. These factors may also manifest themselves as increased variation and explain the relatively wide reference intervals that were observed, although we don't have data to support this.

sOB-R in relation to previous publications

We observed no difference in the sOB-R concentrations between sexes. Nor did we demonstrate an age-dependent correlation with sOB-R concentrations throughout the analyzed age span; however, we did observe significantly elevated sOB-R in the younger children, i.e. girls younger than 10 and boys younger than eight. These findings are also largely in concordance with previous findings [19,40], that demonstrated higher values in younger children, albeit up to 11 years in both sexes. In contrast to the sex-specific leptin pattern, the sOB-R concentrations were inversely correlated to age in both sexes and correspondingly we found the highest sOB-R concentrations in prepubertal children, i.e. Tanner stage I. It has been proposed that the high concentrations of

sOB-R in early childhood may at least in part suppress leptin actions on its membrane receptors [19], but this study does not provide sufficient data to elucidate this theory. We observed a correlation between FLI and leptin (Figure 2), which indicates that the plasma concentration of leptin in healthy, non-obese children is relatively independent of the sOB-R concentration.

Study strengths

Several variables are known to potentially influence leptin measurements [25]. Accordingly, care was taken to minimize the preanalytical [44] variables as all samples were taken in a relatively narrow time span and after an overnight fast which should reduce the influence from the circadian and ultracircadian leptin cycling [45–47] as well as the potential influence from leptin interactions with insulin [48].

As all measurements in this study were done in-house and in batch using the same method, the presented data can be considered valid regarding potential biases, i.e. variation in the analysis methods. The structured family-based questionnaire concerning general health data was self-reported and therefore subject to potential bias. However, it is relatively unlikely for any systematized bias to have influenced the data.

Study weakness

Several statistical methods exist for dealing with interpreting data which include values below LOD, including deletion of non-detects, single substitution, extrapolation by regression, and multiple imputation using TOBIT modeling. We applied the latter as it is believed to be more accurate [34,36]. However, regardless of statistical approach to dealing with values below LOD, data are per definition skewed. Therefore, the sOB-R values $<1.56 \mu\text{g/L}$ must be interpreted with caution. Considering the relatively low percentage of non-detects ($< 10\%$) and the range of sOB-R concentration observed, it is reasonable to presume the influence is small. A potential weakness is pre analysis storage of samples. However, an analysis of the pre-analytic variability demonstrated stability of leptin and sOB-R in the blood sample. Given that the samples were stored at -80°C and only subjected to one freeze-thaw cycle no pre-analytic condition is likely to have influenced the results.

Conclusion

Hyperleptinaemia has been associated with increased risk of the metabolic syndrome and cardiovascular disease [20–22] and it has been proposed that alterations in plasma leptin concentrations and FLI may help evaluating phenotypically non-obese children by detecting those with more unfavorable risk profiles independent of BMI status [49]. The data presented in this study may therefore potentially help in metabolic risk stratification of children; a task that is likely to gain in importance in the near future as a consequence of

the high prevalence of childhood obesity and its associated complications.

Disclosure statement

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

Funding

The study was part of the research activities of the Danish Childhood Obesity Biobank (ClinicalTrials.gov ID NCT00928473) as well as related to TARGET (The impact of our genome on individual treatment response in obese children, Grant no. 0603-00484B) and BIOCHILD (Genetics and system biology of childhood obesity in India and Denmark). The authors wish to thank Mrs Pia Lind and Mrs Oda Troest for expert technical assistance.

References

- Rosenbaum M, Leibel RL. 20 years of leptin: role of leptin in energy homeostasis in humans. *J Endocrinol* 2014;223:T83–96.
- Park H-K, Ahima RS. Physiology of leptin: energy homeostasis, neuroendocrine function and metabolism. *Metab Clin Exp* 2015; 64:24–34.
- Forbes S, Bui S, Robinson BR, Hochgeschwender U, Brennan MB. Integrated control of appetite and fat metabolism by the leptin-proopiomelanocortin pathway. *Proc Natl Acad Sci USA* 2001;98:4233–7.
- Prieur X, Tung YCL, Griffin JL, Farooqi IS, O'Rahilly S, Coll AP. Leptin regulates peripheral lipid metabolism primarily through central effects on food intake. *Endocrinology* 2008;149:5432–9.
- Friedman JM. A tale of two hormones. *Nat Med* 2010;16:1100–6.
- Minokoshi Y, Kim Y-B, Peroni OD, Fryer LGD, Müller C, Carling D, Kahn BB. Leptin stimulates fatty-acid oxidation by activating AMP-activated protein kinase. *Nature* 2002;415:339–43.
- Donoso MA, Muñoz-Calvo MT, Barrios V, Garrido G, Hawkins F, Argente J. Increased circulating adiponectin levels and decreased leptin/soluble leptin receptor ratio throughout puberty in female ballet dancers: association with body composition and the delay in puberty. *Eur J Endocrinol* 2010;162:905–11.
- Agarwal SK, Vogel K, Weitsman SR, Magoffin DA. Leptin antagonizes the insulin-like growth factor-I augmentation of steroidogenesis in granulosa and theca cells of the human ovary. *J Clin Endocrinol Metab* 1999;84:1072–6.
- Matarese G, Procaccini C, De Rosa V, Horvath TL, La Cava A. Regulatory T cells in obesity: the leptin connection. *Trends Mol Med* 2010;16:247–56.
- Sadaf Farooqi I, Matarese G, Lord GM, Keogh JM, Lawrence E, Agwu C, Sanna V, Jebb SA, Perna F, Fontana S, Lechler RI, Depaoli AM, O'Rahilly S. Beneficial effects of leptin on obesity, T cell hyporesponsiveness, and neuroendocrine/metabolic dysfunction of human congenital leptin deficiency. *J Clin Invest* 2002; 110:1093–103.
- Paz-Filho G, Mastrorandi CA, Licinio J. Leptin treatment: facts and expectations. *Metab Clin Exp* 2015;64:146–56.
- Zuo H, Shi Z, Yuan B, Dai Y, Wu G, Hussain A. Association between serum leptin concentrations and insulin resistance: a population-based study from China. *PLoS One* 2013;8:e54615.
- Reinehr T, Kratzsch J, Kiess W, Andler W. Circulating soluble leptin receptor, leptin, and insulin resistance before and after weight loss in obese children. *Int J Obes* 2005;29:1230–5.
- Lönnqvist F, Arner P, Nordfors L, Schalling M. Overexpression of the obese (ob) gene in adipose tissue of human obese subjects. *Nat Med* 1995;1:950–3.
- Huang L, Wang Z, Li C. Modulation of circulating leptin levels by its soluble receptor. *J Biol Chem* 2001;276:6343–9.

16. Schaab M, Kratzsch J. The soluble leptin receptor. *Best Pract Res Clin Endocrinol Metab* 2015;29:661–70.
17. Lammert A, Kiess W, Bottner A, Glasow A, Kratzsch J. Soluble leptin receptor represents the main leptin binding activity in human blood. *Biochem Biophys Res Commun* 2001;283:982–8.
18. Yang G, Ge H, Boucher A, Yu X, Li C. Modulation of direct leptin signaling by soluble leptin receptor. *Mol Endocrinol* 2004;18:1354–62.
19. Kratzsch J, Lammert A, Bottner A, Seidel B, Mueller G, Thierry J, Hebebrand J, Kiess W. Circulating soluble leptin receptor and free leptin index during childhood, puberty, and adolescence. *J Clin Endocrinol Metab* 2002;87:4587–94.
20. Gonzaga NC, Medeiros CC, De Carvalho DF, Alves JG. Leptin and cardiometabolic risk factors in obese children and adolescents. *J Paediatr Child Health* 2014;50:707–12.
21. Yadav A, Kataria MA, Saini V, Yadav A. Role of leptin and adiponectin in insulin resistance. *Clin Chim Acta* 2013;417:80–4.
22. Huang K, Lin RCY, Kormas N, Lee L, Chen C, Gill TP, Caterson ID. Plasma leptin is associated with insulin resistance independent of age, body mass index, fat mass, lipids, and pubertal development in nondiabetic adolescents. *Int J Obes Relat Metab Disord* 2004;28:470–5.
23. Han JC, Lawlor DA, Kimm SYS. Childhood obesity. *Lancet* 2010;375:1737–48.
24. Erhardt E, Foraita R, Pigeot I, Barba G, Veidebaum T, Tornaritis M, Michels N, Eiben G, Ahrens W, Moreno LA, Kovács E, Molnár D. Reference values for leptin and adiponectin in children below the age of 10 based on the IDEFICS cohort. *Int J Obes* 2014;38:S32–S8.
25. Venner AA, Doyle-Baker PK, Lyon ME, Fung TS. A meta-analysis of leptin reference ranges in the healthy paediatric prepubertal population. *Ann Clin Biochem* 2009;46:65–72.
26. Barlow SE. Expert committee recommendations regarding the prevention, assessment, and treatment of child and adolescent overweight and obesity: summary report. *Pediatrics* 2007;120:S164–S92.
27. Cole TJ, Green PJ. Smoothing reference centile curves: the LMS method and penalized likelihood. *Stat Med* 1992;11:1305–19.
28. Nysom K, Mølgaard C, Hutchings B, Michaelsen KF. Body mass index of 0 to 45-y-old Danes: reference values and comparison with published European reference values. *Int J Obes Relat Metab Disord* 2001;25:177–84.
29. Marshall WA, Tanner JM. Variations in pattern of pubertal changes in girls. *Arch Dis Child* 1969;44:291–303.
30. Marshall WA, Tanner JM. Variations in the pattern of pubertal changes in boys. *Arch Dis Child* 1970;45:13–23.
31. van der Sluis IM, de Ridder MAJ, Boot AM, Krenning EP, de Muinck Keizer-Schrama SMPF. Reference data for bone density and body composition measured with dual energy x ray absorptiometry in white children and young adults. *Arch Dis Child* 2002;87:341–7.
32. Yannakoulia M, Yiannakouris N, Blüher S, Matalas AL, Klimis-Zacas D, Mantzoros CS. Body fat mass and macronutrient intake in relation to circulating soluble leptin receptor, free leptin index, adiponectin, and resistin concentrations in healthy humans. *J Clin Endocrinol Metab* 2003;88:1730–6.
33. Rigby RA, Stasinopoulos DM, Lane PW. Generalized additive models for location, scale and shape. *Appl Stat* 2005;54:507–54.
34. Uh H-W, Hartgers FC, Yazdanbakhsh M, Houwing-Duistermaat JJ. Evaluation of regression methods when immunological measurements are constrained by detection limits. *BMC Immunol* 2008;9:59.
35. Hornung RW, Reed LD. Estimation of average concentrations in the presence of nondetectable values. *Appl Occup Environ Hyg* 1990;5:46–51.
36. Andersen A, Benn CS, Jørgensen MJ, Ravn H. Censored correlated cytokine concentrations: multivariate tobit regression using clustered variance estimation. *Stat Med* 2013;32:2859–74.
37. Taylor PJ. Matrix effects: the Achilles heel of quantitative high-performance liquid chromatography-electrospray-tandem mass spectrometry. *Clin Biochem* 2005;38:328–34.
38. Van Eeckhaut A, Lanckmans K, Sarre S, Smolders I, Michotte Y. Validation of bioanalytical LC-MS/MS assays: evaluation of matrix effects. *J Chromatogr B Analyt Technol Biomed Life Sci* 2009;877:2198–207.
39. Sasse EA. Determination of reference intervals in the clinical laboratory using the proposed guideline National Committee for Clinical Laboratory Standards C28-P. *Arch Pathol Lab Med* 1992;116:710–13.
40. Quinton ND, Smith RF, Clayton PE, Gill MS, Shalet S, Justice SK, Simon SA, Walters S, Postel-Vinay MC, Blakemore AIF, Ross RJM. Leptin binding activity changes with age: the link between leptin and puberty. *J Clin Endocrinol Metab* 1999;84:2336–41.
41. Mantzoros CS, Flier JS, Rogol AD. A longitudinal assessment of hormonal and physical alterations during normal puberty in boys. V. Rising leptin levels may signal the onset of puberty. *J Clin Endocrinol Metab* 1997;82:1066–70.
42. Ankarberg-Lindgren C, Dahlgren J, Carlsson B, Rosberg S, Carlsson L, Wikland KA, Norjavaara E. Leptin levels show diurnal variation throughout puberty in healthy children, and follow a gender-specific pattern. *Eur J Endocrinol* 2001;145:43–51.
43. Nagy TR, Gower BA. Correlates of leptin concentration in the San Antonio heart study. *Ann Epidemiol* 1997;7:79–80.
44. Narayanan S. The preanalytic phase. An important component of laboratory medicine. *Am J Clin Pathol* 2000;113:429–52.
45. Saad MF, Riad-Gabriel MG, Khan A, Sharma A, Michael R, Jinagouda SD, Boyadjian R, Steil GM. Diurnal and ultradian rhythmicity of plasma leptin: effects of gender and adiposity. *J Clin Endocrinol Metab* 1998;83:453–9.
46. Sinha MK, Sturis J, Ohannesian J, Magosin S, Stephens T, Heiman ML, Polonsky KS, Caro JF. Ultradian oscillations of leptin secretion in humans. *Biochem Biophys Res Commun* 1996;228:733–8.
47. Licinio J, Negrao AB, Mantzoros C, Kaklamani V, Wong ML, Bongiorno PB, Negro PP, Mulla A, Veldhuis JD, Cernal L, Flier JS, Gold PW. Sex differences in circulating human leptin pulse amplitude: clinical implications. *J Clin Endocrinol Metab* 1998;83:4140–7.
48. Wagner R, Oberste-Berghaus C, Herpertz S, Blum WF, Pelz B, Hebebrand J, Senf W, Mann K, Albers N. Time relationship between circadian variation of serum levels of leptin, insulin and cortisol in healthy subjects. *Horm Res* 2000;54:174–80.
49. Stakos DA, Papaioannou HI, Angelidou I, Mantadakis E, Paraskakis E, Tsigalou C, Chatzimichael A. Plasma leptin and adiponectin concentrations correlate with cardiometabolic risk and systemic inflammation in healthy, non-obese children. *J Pediatr Endocrinol Metab* 2014;27:221–8.